

# Preventing Respiratory Depression

Daniel I. Sessler, M.D.

LEE *et al.*<sup>1</sup> evaluated closed malpractice claims related to respiratory depression. They identified 92 closed claims over 20 yr in about a third of covered anesthesiologists, which corresponds very roughly to 14 closed claims per year for all anesthesiologists nationwide among perhaps 75 million cases. Three-quarters of the patients died or were left with severe brain damage; half resulted in settlement payments, with the median being \$217,000 (intraquartile range: \$50,000 to \$604,000). We know that only a small fraction of adverse outcomes results in malpractice claims. It is thus apparent that postoperative respiratory events resulting in death or serious injury occur at a concerning rate.

Most patients whose closed claims resulted from respiratory events were given opioids, nearly half by at least two routes—often prescribed by different physicians, and nearly half had a continuous opioid infusion at the time of the respiratory event. That said, gross overdose apparently contributed to only 16% of cases. Interestingly, only a quarter of the patients had or were at high risk of obstructive sleep apnea, which is consistent with data showing no relationship whatsoever between STOP-BANG scores (a measure of sleep apnea risk)<sup>2</sup> and postoperative hypoxemia.<sup>3</sup>

Taken together, these data suggest that routine opioid use is a major cause of postoperative respiratory events—probably contributing more than sleep apnea. Reducing opioid use seems likely to decrease the risk of respiratory events. However, it is also obvious that opioids will remain an important analgesic strategy for years to come—and that we must thus learn to safely care for patients given opioids.

Postoperative airway obstruction or inadequate ventilatory effort should not cause catastrophic outcomes because either can be treated with opioid antagonists or mechanical ventilation. Respiratory catastrophes after routine surgery might thus reasonably be considered “never events.”



***“It is likely that many catastrophic respiratory events could be prevented by continuous ... monitoring. However, major trials are needed to determine what should be monitored and how.”***

Case reviewers judged that 97% of claims probably or possibly could have been prevented by better monitoring. More intense conventional monitoring probably is not the answer though. Hypoxemia in postoperative inpatients is common, severe, and prolonged.<sup>4</sup> Furthermore, even serious and persistent hypoxemia is unrecognized by nurses in 88% of cases (unpublished data, Department of Outcomes Research, Cleveland Clinic, December 2014). That a full quarter of the respiratory events in the series of Lee *et al.*<sup>1</sup> occurred within 15 min of a nursing evaluation shows how often the current system fails.

So what can we do? Continuous monitoring is perhaps the obvious way to prevent catastrophic postoperative respiratory events. It is tempting to target continuous monitoring to high-risk patients, such as those who have a history of sleep apnea or are obese. The difficulty is that even the best prediction system will identify only a fraction—probably a small fraction—of at-risk patients. Continuous monitoring is thus probably appropriate for nearly all postoperative inpatients.

The real question is perhaps not *whether* to monitor continuously, but *what* to monitor. Surely pulse oximetry should be included. But whether to add capnography, impedance pneumography, transcutaneous partial pressure of carbon dioxide, or other monitoring systems is less obvious. (The Anesthesia Patient Safety Foundation recommends pulse oximetry for all patients given parenteral opioids and some measure of ventilation for patients using supplemental oxygen.)<sup>5</sup> Clearly, each type of ventilation monitoring provides valuable information that supplements oxygen saturation, but each also adds complexity, cost, and false alarms. Recently developed compact battery-powered systems that communicate wirelessly reduce the burden for patients. And as interpretation algorithms become more sophisticated, the ratio of true-positive to false-positive alarms will presumably improve.

Image: ©Thinkstock.

Corresponding article on page 659.

Accepted for publication November 4, 2014. From the Department of Outcomes Research, Cleveland Clinic, Cleveland, Ohio.

Copyright © 2014, the American Society of Anesthesiologists, Inc. Wolters Kluwer Health, Inc. All Rights Reserved. *Anesthesiology* 2015; 122:484-5

Nonetheless, the history of medicine is littered with treatments that seemed logical and were subsequently proven unhelpful—or even harmful. There is little reason to believe that our understanding of physiology has suddenly become so good that logic alone should suffice in this case. Continuous respiratory monitoring is no different from any other clinical interventions and deserves similar formal testing including a cost-benefit analysis. The question is not simply whether to monitor; we also need to know which patients, what system(s) to use, and how to interpret the resulting data.

In summary, half the closed claims postoperative respiratory events were lethal and a quarter caused serious neurologic injury. Opioid administration probably contributed in most cases, and inadequate monitoring was the rule. It is likely that many catastrophic respiratory events could be prevented by continuous saturation and/or ventilation monitoring. However, major trials are needed to determine what should be monitored and how.

### Acknowledgments

The author has received funding from Covidien (Dublin, Ireland) for a study of postoperative hypoxemia. He has no personal financial interest associated with the topic of this article.

### Competing Interests

Sotera Wireless (San Diego, California), a company that makes a continuous postoperative monitor, donat-

ed equipment for a Department of Outcomes Research (Cleveland Clinic, Cleveland, Ohio) study. None of the results cited in this editorial were obtained with Sotera equipment.

### Correspondence

Address correspondence to Dr. Sessler: ds@or.org

### References

1. Lee LA, Caplan RA, Stephens LS, Posner KL, Terman GW, Voepel-Lewis T, Domino KB: Postoperative opioid-induced respiratory depression: A closed claims analysis. *ANESTHESIOLOGY* 2015; 122:659–65
2. Farney RJ, Walker BS, Farney RM, Snow GL, Walker JM: The STOP-Bang equivalent model and prediction of severity of obstructive sleep apnea: Relation to polysomnographic measurements of the apnea/hypopnea index. *J Clin Sleep Med* 2011; 7:459–65B
3. Khanna AK, Sun Z, Naylor A, You J, Hesler BD, Kurz AM, Devereaux PJ, Sessler DI, Saager L: The STOP-BANG questionnaire as a predictor of hypoxia in non-cardiac surgery patients: A prospective cohort analysis. Presented at: Annual Meeting of American Society of Anesthesiologists, New Orleans, Louisiana, October 14, 2014 (A4057)
4. Shahinyan A, Sun Z, Finnegan PS, Egan C, Fu A, Devereaux PJ, Dalton J, Sessler DI, Kurz A: Duration and incidence of postoperative hypoxemia. Paper presented at: Annual Meeting of American Society of Anesthesiologists, San Francisco, California, October 15, 2013 (BOC12)
5. Weigner MB, Lee LA: No patient shall be harmed by opioid-induced respiratory depression. *APSF Newsletter* 2011; 26:21, 26–8